

Lipopolysaccharide binding protein (L.B.P.) – an inflammatory marker of prognosis in the acute appendicitis

Brănescu C*, Șerban D**, Șavlovschi C**, Dascălu AM**, Kraft A**

*IVth Surgery Clinic, University Emergency Hospital, Bucharest, Romania

**"Carol Davila" University of Medicine and Pharmacy, Bucharest, Romania

Correspondence to: Cristian Brănescu, Attending surgeon, Doctoral candidate

University Emergency Hospital, Bucharest, Romania,

169 Splaiul Independentei, RO – 050098, Bucharest, Romania

4th Surgery Clinic, E-mail: cristi.branescu@gmail.com

Received: June 1st, 2012 – Accepted: September 6th, 2012

Abstract

LIPOPOLYSACCHARIDE BINDING PROTEIN (LBP) is an important mediator of the inflammatory reaction. A multitude of factors can determine the genic transcription activation and the increase of the LBP in the blood and the human body humours: IL1, IL6, lipopolysaccharides, Gram-negative bacteria, as well as non-infectious agents.

This paper is a prospective study performed on 147 patients admitted for acute appendicitis in 2010-2012 and evaluates the dynamics of LBP in acute appendicitis, by identifying the correlations between the pre- and post-operative levels of LBP (up to 72 hours after surgery) and the anatomopathological type (i.e. catarrhal, phlegmonous and gangrenous).

The mean pre-op LBP values are significantly different as to the histopathological result ($p < 0,005$). Among the biological inflammatory markers measured in this present study, LBP has a dynamics of its own in the catarrhal and phlegmonous appendicitis. Thus, if after the surgical removal of the infectious source, the leukocytes and neutrophiles decrease 72 hrs after surgery, LBP continues an ascending curve.

The importance of this study consists in the introduction of last generation LBP-type inflammatory markers' dosage in the cecal appendix pathology. This implementation is brand new in the Romanian surgical practice. The good correlation between the LBP pre-op values and the histopathological diagnosis of the appendicitis form that we discovered during the present study opens the way to large-scale use of the biochemical dosage of LBP in the management of acute appendicitis.

- **Keywords:** Acute appendicitis, Lipopolysaccharide binding protein (LBP), anatomopathology, acute phase proteins •

Abbreviations: Lipopolysaccharide binding protein – LBP; Anatomopathology – AP

Introduction

The first appendicitis performed on December 6th 1735 [1] by Claudius Amyand was introduced as therapeutically relevant for Amyand's Hernia [2]. In time, the science has made significant progress and the inflammatory markers' era has begun to identify increasingly detailed sequences of each stage of the surgical pathology.

LBP is a 50-kDa polypeptide mainly synthesized in hepatocytes and is released as a 58- to 60-kDa glycoprotein into the bloodstream after glycosylation [3]. The LPS-binding domain has been identified in the N-terminal end of LBP comparable to the structural and functional related bactericidal permeability increasing protein (BPI) [4]. Both the LBP gene and the BPI gene are closely located on the chromosome 20 [5]. The transcriptional regulation of the LBP gene is induced by interleukin (IL)-1 alone or synergistically by IL-1 and IL-6, leading to a maximal LBP concentration within 24-48 h after stimulation, a response that can be strongly enhanced by tumor necrosis factor (TNF)-α and dexamethasone [6]. Other stimuli inducing LBP-synthesis

in vivo include LPS, Gram-negative bacteria, and non-infectious agents such as turpentine [7]. The hepatic transcriptional induction of LBP is inhibited by the anti-inflammatory cytokine transforming growth factor β1 (TGF-β1) [8].

LBP is constitutively present in serum at concentrations of 5e15 mg/ml and rises 10- to 50-fold during the acute-phase reaction [9].

The aim of present article is to identify the role played by inflammatory markers as prognosis factors in the acute appendicitis, mainly that of LBP; the LBP monitorization and quantification in the diagnosis stage is meant to confirm the choice of the proper surgical conduct.

Materials and method

The aim of the study was to assess, based on the available preoperative information, the following aspects:

- ▲ the accuracy of the intra-operative result prediction;
- ▲ the evolution of some parameters between the admission moment and the analysis of the correlations and associations among the pre-operative factors.

A prospective study was therefore performed, based on a sample of 147 cases of acute appendicitis operated in 2010-2012, in the 4th Surgery Clinic of the University Emergency Hospital, Bucharest.

The pre-operative information consisted in:

- demographic data (i.e. age, sex, place of origin);
- clinical signs (i.e. spontaneous pain or upon palpation, tenderness, muscular defense, nausea, vomiting, bowel movement disorders, fever);
- lab tests (i.e. leukocytes, leukocyte formula, LBP).

The intra-operative result the surgeon settled by direct observation during the intervention consisted in assigning each patient to one of the following appendicitis classes: catarrhal, phlegmonous, gangrenous, with possible additional clarifications regarding the occurrence of peritonitis or perforation. The anatomopathological

diagnosis was decided based on the histopathological exam which divided the excision specimens into three groups, as follows: catarrhal, phlegmonous and gangrenous.

Those susceptible of having a second infection source – other than the one caused by appendicitis – were expelled. The rationale for the expel criteria is that we intended to seize the dynamics of change of the biological inflammatory markers since the admission moment up to 72 hrs after surgery, on condition that the unique infection source – caused by appendicitis – be eliminated.

Results


Out of the 147 patients included in the sample, 79 (53.7%) were females and 68(46.3%) males. 54.4% were catarrhal appendicitis, 32.7% phlegmonous and 12.9% gangrenous (**Table 1**).

Table 1. Distribution of anatomopathological result on sexes

Sex	F	Anatomopathological Result			Total
		Catarrhal	Phlegmonous	Gangrenous	
%	F	54	22	3	79
	%	68.4%	27.8%	3.8%	100.0%
M	Nr	26	26	16	68
	%	38.2%	38.2%	23.5%	100.0%
Total	Nr	80	48	19	147
	%	54.4%	32.7%	12.9%	100.0%

The distribution of catarrhal, phlegmonous and gangrenous appendicitis differs significantly statisticwise with SEX ($p<0.001$; Pearson's test χ^2). It is noteworthy that with female patients 68% of the cases are catarrhal appendicitis, and only 3.8% are gangrenous, while with male patients the percentages are different: there are significantly more cases of gangrenous appendicitis and less catarrhal ones (23.5% and 38.2%, respectively) (**Graph 1**).

Graph 1. Distribution of anatomopathological result on sexes




The age distribution (see **Table 2**) differs significantly with the AP Result ($p<0.001$; the ANOVA test). It is easy to notice that the result "gangrenous appendicitis" usually associates with higher ages. Concretely, the average age of the patients is maximum in the case of gangrenous appendicitis, whereas for the catarrhal and phlegmonous ones it is 14-15 years as low.

Table 2. Average age as to the AP Result

AP Result	Mean	N	Std. Deviation	Median
Catarrhal	27.95	80	8.846	26.00
Phlegmonous	28.83	48	12.931	24.00
Gangrenous	43.32	19	16.954	38.00
Total	30.22	147	12.545	26.00

Graph 2 leads to the same conclusions, but highlights a different aspect, i.e. the majority of patients with catarrhal and phlegmonous appendicitis are aged between 18-30, whereas the patients with gangrenous ones are relatively evenly distributed between 20-60 years of age.

Graph 2. Age distribution (frequency) as to the Anatomopathological Result



Graph 3 shows the distribution of the anatomopathological result as to the age groups 18-30, 31-50 and over 51.

Graph 3. Distribution of the anatomopathological result as to the age groups






Table 3. Preoperative LBP, number of Leukocytes, Neutrophiles and Alvarado Score are significantly different, as to the acute appendicitis type inflammation. The increasing order of the average values is the same for all laboratory tests: Catarrhal < Phlegmonous < Gangrenous.

The only exception from the previous statement refers to Neutrophiles. Neutrophiles' Frequency is significantly lower in the case of catarrhal appendicitis than in the case of phlegmonous and gangrenous ones, but the values for these last two forms does not differ significantly.

Leukocytosis with neutrophilia occurs in 100% of acute gangrenous appendicitis (**Table 3**).

Table 3. Preoperative LBP, number of Leukocytes, Neutrophiles and Alvarado Score

AP Result		Preop LBP	No. of Leukocytes upon admission	Leuko-cytosis 1000	Neutro-philia	Alvarado Score
Catarrhal (80)	Average	1.937	8908.00	0.25	0.27	6.55
	Std. Dev.	1.2909	3200.814	0.44	0.45	0.79
	Median	1.600	8455.00	0	0	7.00
	Minimum	0.4	3900	0	0	5
	Maximum	4.9	21980	1	1	8
Phlegmonous (48)	Average	5.583	13622.92	0.87	0.96	8.65
	Std. Dev.	1.7040	3607.966	0.33	0.2	0.7
	Median	5.500	12950.00	1.00	1.00	9.00
	Minimum	3.1	7500	0	0	7
	Maximum	8.9	21250	1	1	10
Gangrenous (19)	Average	33.826	18114.74	1.00	1.00	9.63
	Std. Dev.	5.0224	3450.276	0	0	0.6
	Median	35.700	17500.00	1.00	1.00	10.00
	Minimum	20.9	10680	1	1	8
	Maximum	39.1	23100	1	1	10
Total (147)	Average	7.250	11637.55	0.55	0.59	7.63
	Std. Dev.	10.6414	4694.671	0.5	0.49	1.429
	Median	3.700	10500.00	1.00	1.00	7.00
	Minimum	0.4	3900	0	0	5
	Maximum	39.1	23100	1	1	10

Table 4. Shows the main statistical indicators for the post-op laboratory tests. The following conclusions can be drawn:

The mean values of LBP, Leukocyte number are significantly different, with the acute appendicitis type inflammation. The increase order of the mean values is the same for all tests: Catarrhal < Phlegmonous < Gangrenous;

The only exception to the previous statement refers to LBP: unlike the difference between the means which was very big pre-op, in post-op this difference becomes insignificant between the phlegmonous and the gangrenous form.

Table 4. Main statistical indicators for the post-op laboratory tests

AP Result	LBP Postop	No. of Leucocytes after 72 hrs
Cataral (80)	Mean	7,446
	Std. Deviation	1,1861
	Median	7,400
	Minimum	5,1
	Maximum	9,9
Flegmonos	Mean	12,460
		8541,04

	Std. Deviation	2,2964	2309,748
	Median	12,600	8100,00
	Minimum	6,5	4700
Gangrenos	Maximum	15,9	13600
	Mean	12,547	11417,89
	Std. Deviation	2,8541	2068,546
	Median	12,400	11500,00
	Minimum	7,1	7700
Total (147)	Maximum	17,5	14600
	Mean	9,743	7575,03
	Std. Deviation	3,1313	2821,643
	Median	8,800	6700,00
	Minimum	5,1	3500
	Maximum	17,5	14600

Table 5. Unidimensional statistical analysis of the pre- and post-op evolution of the biological inflammatory markers with patients in the study sample as to the anatomo-pathological type of appendicitis (Catarrhal, Phlegmonous, Gangrenous) (**Table 5**).



The values of LBP and Leukocytes Number change significantly between pre-op and post-op ($p<0,001$; t test). This can be seen for all 3 anatomo-pathological types (**Table 5**).

AP Result	Variable: Postop-Preop Difference	Test Value = 0				
					95% Confidence Interval of the Difference	
		Mean Difference	t	Sig. (2-tailed)	Lower	Upper
Catarrhal	Dif. LBP	5,50875	28,192	,00	5,1198	5,8977
	Dif.No.Leukocytes	-2825,25000	-16,260	,00	-3171,1028	-2479,3972
Phlegmonous	Dif.LBP	6,87708	15,332	,00	5,9748	7,7794
	Dif.No.Leukocytes	-5081,87500	-18,930	,00	-5621,9463	-4541,8037
Gangrenous	Dif.LBP	-21,27895	-23,367	,00	-23,1921	-19,3658
	Dif.Nr.Leukocytes	-6696,84211	-14,635	,00	-7658,2239	-5735,4603

The values of the Leukocytes Number decrease significantly after surgery (**Graph 4**).

Mean values of the number of leukocytes upon admission and 72 hrs after surgery


Graph 4. Number of leukocytes evolution between admission value and 72 hrs after surgery




Number of leukocytes: "boxplot" representation of the difference between the value 72 hrs after surgery and the value upon admission

Also after surgery, the LBP values increase significantly with the catarrhal and phlegmonous types, but decrease significantly with the gangrenous one (**Graph 5**).

Graph 5. LBP evolution between admission and post-op values, for the three appendicitis types



Mean LBP values, Pre- and Post-op



LBP: "boxplot" representation of the difference between the pre- and post-op values

Discussion

LBP is a biological marker with high prognostic value in the acute appendicitis. LBP dosage pre- and post-op is useful for discrimination among the three types of appendicitis; its recovery to the range statistically considered as normal confirms the infectious process in the organism was solved.

An interesting remark is that among the biological inflammatory markers dosed in the present study, LBP has a dynamics of its own in catarrhal and phlegmonous appendicitis. Thus, if after the surgical removal of the infection source the leukocytes and neutrophiles values decrease 72 hrs post surgery, LBP continues an ascending curve. A statistically significant decrease of post-op LBP is seen though in the cases of gangrenous appendicitis (in which the pre-op values were significantly higher).

An explanation to this particular response mechanism is the LBP production mechanism: i.e. by the genic transcription adjustment of the IL1 and IL6-mediated proteic synthesis. The LBP synthesis might be not only connected to the infectious factor's presence, but also to the level of inflammation in the human body.

Another possibility might be that the surgical act itself and the post-op wound healing, could stimulate the LBP synthesis by increasing the level of the specific inflammatory mediators, although this stimulation does not reach the extremely increased level seen in the case of gangrenous appendicitis.

The importance of the present study consists in the introduction of the last generation LBP-type inflammatory markers in the cecal appendix pathology, an extremely frequent occurrence in the emergency surgical pathology, but whose clinical-paraclinical picture is polymorphic, raising doubts in the pre-op diagnosis – especially at the onset of its evolution. This implementation is brand new in the Romanian surgical practise.

The good correlation between the LBP pre- and post-op values and the hystopathological diagnosis of the appendicitis form we have identified and presented in this study gives the opportunity to use LBP biochemical dosage on a large scale in the management of acute appendicitis.

Disclosure: No financial interests

References

- Milanchi S, Allins A D. Amyand's hernia: history, imaging and management. *Hernia*. 2008;12:321-322.
- Şavlovschi C, Brănescu C, Şerban D, Tudor C, Găvan C, Shanabli A, Comandaşu M, Vasilescu L, Borcan R, Dumitrescu D, Sandolache B, Sajin M, Grădinaru S, Munteanu R, Kraft A, Oprescu S. Hernia Amyand – caz clinic. *Chirurgia*. 2010;(3)105: 409-414.
- Zweigner J, Schumann R.R, Weber JR. The role of lipopolysaccharide-binding protein in modulating the innate immune response. *Microbes and Infection* 8.2006;946-952.
- Dentener MA, Vreugdenhil AC, Hoet PH, Verwooy JH, Nieman FH, Heumann D, Janssen YM, Buurman WA, Wouters EF. Production of the acute-phase protein lipopolysaccharidebinding protein by respiratory type II epithelial cells: implications for local defense to

- bacterial endotoxins. Am. J. Respir. Cell Mol. Biol. 23.2000;146e153.
5. **Gray PW, Corcoran AE, Eddy RL Jr., Byers MG, Shows TB.** The genes for the lipopolysaccharide binding protein (LBP) and the bactericidal permeability increasing protein (BPI) are encoded in the same region of human chromosome 20. Genomics 15.1993;188e190.
6. **Geller DA, Kispert PH, Su GL, Wang SC, Di Silvio M, Twardy DJ, Billiar TR, Simmons RL.** Induction of hepatocyte lipopolysaccharide binding protein in models of sepsis and the acute-phase response. Arch. Surg. 128.1993;22e27.
7. **Schumann RR, Kirschning CJ, Unbehau A, Aberle HP, Knope HP, Lamping N, Ulevitch RJ, Herrmann F.** The lipopolysaccharide-binding protein is a secretory class 1 acute-phase protein whose gene is transcriptionally activated by APRF/STAT3 and other cytokine-inducible nuclear proteins. Mol. Cell. Biol. 16.1996;3490e3503.
8. **Hallatschek W, Fiedler G, Kirschning CJ, Creutzburg F, Lamping N, Nussler A, Schumann RR.** Inhibition of hepatic transcriptional induction of lipopolysaccharide-binding protein by transforming-growth-factor beta 1. Eur. J. Immunol. 34.2004;1441e1450.
9. **Zweigner J, Gramm HJ, Singer OC, Wegscheider K, Schumann RR.** High concentrations of lipopolysaccharide-binding protein in serum of patients with severe sepsis or septic shock inhibit the lipopolysaccharide response in human monocytes. Blood. 98.2001;3800e3808.